

Jack, C., Hotchkiss, E., Sargison, N. D., Toma, L., Milne, C. and Bartley, D. J. (2017) A quantitative analysis of attitudes and behaviours concerning sustainable parasite control practices from Scottish sheep farmers. *Preventive Veterinary Medicine*, 139, pp. 134-145.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/201457/>

Deposited on: 23 October 2019

A quantitative analysis of attitudes and behaviours concerning sustainable parasite control
practices from Scottish sheep farmers

Corin Jack ^{a,*}, Emily Hotchkiss ^a, Neil D. Sargison ^b, Luiza Toma ^c, Catherine Milne ^c, David.
J. Bartley ^a

^a Disease control, Moredun Research Institute, Pentlands Science Park, Bush Loan, Penicuik,
EH26 0PZ, UK.

^b Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush Campus,
Midlothian, EH25 9RG, UK

^c Scotland's Rural College (SRUC), Peter Wilson Building, Kings Buildings, West Mains
Road, Edinburgh EH9 3JG, UK

*Corresponding author. Tel.: +44 0131 4455111; fax: +44 0131 4455222

E-mail address: corin.jack@moredun.ac.uk

Keywords: Behaviour; Parasite control; Questionnaire; S.E.M; Sheep; *Structural Equation
Modelling*

Abstract

Nematode control in sheep, by strategic use of anthelmintics, is threatened by the emergence of roundworms populations that are resistant to one or more of the currently available drugs. In response to growing concerns of Anthelmintic Resistance (AR) development in UK sheep flocks, the Sustainable Control of Parasites in Sheep (SCOPS) initiative was set up in 2003 in order to promote practical guidelines for producers and advisors. To facilitate the uptake of 'best practice' approaches to nematode management, a comprehensive understanding of the various factors influencing sheep farmers' adoption of the SCOPS principles is required.

A telephone survey of 400 Scottish sheep farmers was conducted to elicit attitudes regarding roundworm control, AR and 'best practice' recommendations. A quantitative statistical analysis approach using structural equation modelling was chosen to test the relationships between both observed and latent variables relating to general roundworm control beliefs. A model framework was developed to test the influence of socio-psychological factors on the uptake of sustainable (SCOPS) and known unsustainable (AR selective) roundworm control practices. The analysis identified eleven factors with significant influences on the adoption of SCOPS recommended practices and AR selective practices. Two models established a good fit with the observed data with each model explaining 54% and 47% of the variance in SCOPS and AR selective behaviours, respectively. The key influences toward the adoption of best practice parasite management, as well as demonstrating negative influences on employing AR selective practices were farmer's base line understanding about roundworm control and confirmation about lack of anthelmintic efficacy in a flock. The findings suggest that improving farmers' acceptance and uptake of diagnostic testing and improving underlying knowledge and awareness about nematode control may influence adoption of best practice behaviour.

47

48 **Introduction**

49 The sustainable control of gastro-intestinal nematode parasites remains one of the main
50 perennial endemic disease pressures that livestock farmers face globally (Jackson and Coop,
51 2000; Nieuwhof and Bishop, 2005). Gastro-intestinal nematodes impact on the health,
52 welfare and production efficiency of livestock (Coop and Kyriazakis, 2001). For over 50
53 years parasite control strategies have heavily relied on suppressing nematode populations
54 with frequent use of highly efficacious, broad spectrum anthelmintics (Bartley, 2008). The
55 effectiveness of these treatments is threatened by the emergence of nematode populations that
56 are resistant to one or more of the anthelmintic drugs available. In the UK alone, studies have
57 reported resistance to all three of the commercially available broad-spectrum anthelmintic
58 drug classes i.e. benzimidazoles (1-BZ), levamisole (2-LV) and macrocyclic lactones (3-
59 ML). Widespread 1-BZ resistance has been reported throughout the UK (Cawthorne and
60 Whitehead, 1983; Sutherland et al., 1988; Grimshaw et al., 1994; Bartley et al., 2003;
61 Mitchell et al., 2010; Thomas et al., 2015), with a much lower number of 2-LV resistance
62 reports observed (Hong et al., 1994; Coles and Simkins, 1996; Mitchell et al., 2010) and
63 increasing reports of 3-ML resistance associated with multiple drug resistance to two or more
64 anthelmintic drug classes (Bartley et al., 2004; Sargison et al., 2005; Sargison et al., 2007;
65 Thomas et al., 2015). It is therefore increasingly apparent that taking steps toward
66 maintaining sustainable productivity in the growing face of anthelmintic resistance (AR) is
67 required by farmers.

68 In response to growing concerns of AR development in the UK sheep industry, the
69 Sustainable Control of Parasites in Sheep (SCOPS) initiative was set up in 2003. SCOPS is
70 an industry led group that represents the interests of the UK sheep industry with a remit to

develop and promote practical recommendations for producers and advisors regarding ‘best practice’ approaches to parasite control (Abbott et al., 2012). Currently these recommendations are summarised into eight guidelines each of which outline a variety of measures to preserve the effectiveness of current and future anthelmintics. These eight guidelines broadly cover the following aspects of best practice roundworm control including:

- 1) Working out a control strategy with a veterinary advisor
- 2) implementing an effective quarantine strategy
- 3) testing for anthelmintic resistance,
- 4) administering anthelmintics effectively
- 5) using anthelmintics only when necessary
- 6) selecting the appropriate anthelmintics
- 7) preserving a susceptible worm population and
- 8) introducing alternative, non-chemotherapeutic roundworm control strategies (Abbott et al., 2012).

There are numerous channels for the dissemination of the SCOPS recommendations such as through animal health advisors (e.g. veterinarians, suitably qualified persons and researchers), online/printed publications as well as face-to-face promotion at agricultural events. In other sheep producing countries such as Australia, the current equivalent repository for information and recommended practices regarding roundworm control *WormBoss* (Anonymous, 2016) has achieved a high level of awareness amongst farmers. This is in part due to the effective use of the internet platform including the use of an electronic support system. However steps to measure and enhance the transition from awareness to adoption are an uncertainty recognised by both extension schemes (Woodgate and Love, 2012; Anonymous, 2013)).

Various questionnaire surveys have been undertaken and published on the parasite management practices of sheep farmers from around the world, as well as within the UK (Coles, 1997; Bartley et al., 2004; Suter et al., 2005; Hughes et al., 2007; Lawrence et al., 2007; Sargison et al., 2007; Morgan et al., 2012; McMahon et al., 2013). Such studies have highlighted the variable adoption of sustainable roundworm control practices, and emphasised the need to improve promotion and perception of these practices if sustainable

parasite control is to be generally accepted. In recent years the rapidly growing application of socio-psychological research methods in behavioural science has highlighted their influence on animal health decision making. These studies have investigated behaviours relating to a wide range of disease management practices related to many livestock species as described by Wauters & Rojo-Gimeno (2014). However, a limited amount of work has investigated how socio-psychological factors may influence farmer's parasite control behaviours (e.g. Relf et al., 2012; Vande Velde et al., 2015). Moreover few studies have employed the use of quantitative modelling techniques to assess the extent at which such factors influence farmers' parasite control behaviours. The measure of human behaviour in these studies has either been indicated via behavioural intentions (e.g. Toma et al., 2015; Vande Velde et al., 2015) or by respondents' self-reported behaviours (Toma et al., 2013). The use of behavioural intention i.e. a readiness to perform a given behaviour has been proposed to be a direct proxy for actual behaviour based on the widely applied theory of planned behaviour model (Ajzen, 1991). Self-reported behaviour on the other hand requires respondents to personally state their actions regarding a specific circumstance. More recent applications of decision-making models have moved from primarily economic driven factors to also incorporate non-economic influences such as farm characteristics, farmer demographics and psychological factors. This helps to represent the range of both financial and non-financial factors involved and their potential influences in the decision making process (Edwards-Jones, 2006).

This study aims to use a quantitative statistical modelling approach to investigate the influence of socio-psychological factors on the overall adoption of SCOPS practices and practices recognised to be selective for the development of AR (designated AR selective practices hereafter). By employing such methods this will help to evaluate potential

mitigation strategies to assist the adoption of best practice parasite management approaches.

Material and methods

Model framework

Attitudinal questionnaire items were initially devised based on a range of different source material. Questions came from a combination of common themes highlighted from farmer focus group meetings (unpublished data), as well as the research groups own parasite management experience and comparable questionnaire literature related to disease management (Bartley et al., 2003; Palmer, 2009; Toma et al., 2013; Alarcon et al., 2014; Vande Velde et al., 2015). The emphasis for developing questions was to consider areas of greatest importance to sheep farmers regarding parasite control, such as treatment timings, benefits of anthelmintic treatments, dosing practice etc. The result of this was a comprehensive list of items which were categorised into components based on the SCOPS guidelines. Questions that were not specific to SCOPS practices were grouped under ‘general attitudes’ to roundworm control and anthelmintic resistance. Additional items were derived from behavioural models such as the Health belief model (HBM) which has been used to explain and predict preventive health behaviours (Rosenstock et al., 1988). Such items derived from this model include perceived level of risk, which comprises of susceptibility i.e. likelihood of an event occurring, as well as severity i.e. the impact of the event occurring. The combination of these risk items is referred to as ‘risk perception’ and was incorporated into the general attitudes section of the questionnaire. Figure 1 illustrates the model framework used in this paper to examine the influence of general roundworm control and AR

attitudes and farming demographic influences on the overall uptake of SCOPS and known AR selective practices.




Fig.1. Theoretical framework for general uptake of SCOPS recommended and AR selective roundworm control practices

Quantitative attitudinal survey design

The survey design was informed from the model framework (Figure 1) and built around four main components which were arranged in the following order; 1) farmer demographics and enterprise characteristics, 2) general roundworm control/AR attitude statements, 3) open-ended roundworm control knowledge questions and 4) parasite control behaviours. The first section included ten closed-ended questions relating to demographical information (age, education and years earning a living as a farmer), as well as details of the farming system

(e.g. enterprise type, flock size, land topography, farming priorities). The second section included 20 broader questions relating to attitudes towards general parasite control that were not specific toward a particular control measure (e.g. the perceived importance of roundworms, Attitudes to veterinary service and risk perception of AR). The third section included three open-ended questions which were used to gauge the level of the respondent's knowledge and understanding on the topic of roundworm control and AR. The final section included 19 closed-ended questions of which 15 were directed to parasite control measures implemented on farm. Four additional questions which included directly relating to the behaviours of interest as well as preferred formats of knowledge transfer. All attitudinal items included in section 2 were measured on a 5-point-Likert scale: Strongly Disagree (1), Disagree (2), Unsure (3), Agree (4) and Strongly Agree (5). Sections 1 and 4 were recorded by interviewers based on a pre-determined coding frame.

Survey implementation

Farmer contact details were obtained from the Scottish Government (Rural and Environment Science and Analytical Services Division; RESAS) by the use of a stratified simple random sampling method applied to the agricultural census data. The selection criteria used to target farms of interest included, premises with flocks with more than 50 breeding ewes and other sheep (1-year-old and over) for breeding, and at least 25 ewes used for breeding in the previous season. This was to avoid sampling from particular smallholdings where the motives for rearing livestock are not financially driven. The sampling frame was further stratified regionally by animal health divisional office (AHDO) in order to ensure a proportional population sample from each region. Based on a target of 400 completed surveys from across six geographic regions of Scotland, the number required per region was weighted based on the overall number of holdings within the region. The 400 target was established based on a calculated sample size using the number of Scottish sheep holdings (approx.

14,900; National statistics) with an error rate of 5% and confidence level of 95% (Israel, 1992). The following equation was used to calculate the sample size for the questionnaire where n is the sample size, N is the population size, and e is the level of precision (Yamane, 1967)

$$n = \frac{N}{1 + N(e)^2}$$

Opt-out letters were sent out to farmers two weeks prior to the implementation of the survey. The letters outlined the aim of the study, the estimated interview duration, the voluntary nature of the survey and gave assurance that any publication of results would ensure anonymity. If the recipient did not reply to the opt-out letter within the specified time it was considered that they had implicitly agreed to participate in the telephone interview.

A pilot study with six farmers was conducted before undertaking the main survey. This informed the modification of questionnaire items ensuring no ambiguity of questions by respondents and suitability of items for the telephone survey format. Additionally, lengths of interviews were monitored to ensure that interview times were not excessive, in order to achieve appropriate timeliness.

The survey interviews were conducted by a telecommunications company (Feedback Market Research Ltd.) and responses were documented by the interviewer and compiled on a Microsoft Excel spreadsheet. All interviews were conducted under internal quality assurance procedures using computer assisted telephone interviewing systems. Farmers were assured that all information provided would remain completely anonymous in any subsequent reports or publications and that they and their enterprises would not be individually identifiable. Any farmers wishing to opt out after the data was collected were able to do so.

Data formatting/transformation

218 The raw data was firstly coded into a database using Statistical Package for the Social
219 Sciences (SPSS, IBM SPSS Statistics version 22.0). The data was then assessed for normality
220 using the Kolmogorov-Smirnov statistical test. All variables included in the analysis were
221 recorded as per the original coding frame detailed in Table 2, with the exception of
222 ‘Education’, ‘Ewe numbers’ and ‘Roundworm control knowledge’. Categories other than
223 ‘agricultural college’ within ‘Education’ were considered to have little influence on
224 agricultural practice and were therefore combined. The continuous variable ‘Ewe numbers’
225 was categorised based on an evaluation of the data structure. The three open-ended
226 knowledge question responses were individually assessed and classified into a dichotomous
227 variable (i.e. correct or incorrect) based on the authors’ judgement. Two of the three
228 questions required the respondent to list specific examples of parasite species or roundworm
229 control practices. The third question required a description of their understanding of the term
230 wormer resistance, a correct response required a description of the basic principle i.e. a
231 reduction in the effectiveness of a drug treatment or an inherent ability of parasites to survive
232 drug treatment. A score was devised based on the number of correct responses to the three
233 questions.

234 The endogenous i.e. dependent variables (‘SCOPS practice uptake’ & ‘AR selective practice
235 uptake’) were formulated into ordinal scores by summing the total number of practices that
236 were identified as either ‘best practice’ or selective for AR development based on the SCOPS
237 manual (Abbot et al., 2012). The designation of AR selective practices was based on the
238 selection of behaviours which were converse to best practice approaches, also which were
239 impartial towards particular farming systems. The total number of practices identified as best
240 practice was ten, and the total number of AR selective practices identified was seven. Table 1
241 presents the descriptive statistics of the practices used to formulate both outcome variables

(i.e. SCOPS and AR selective practices) and figure 2 presents the frequency distributions of the outcome variables.

Table 1.

Respondents roundworm control behaviours associated with ‘SCOPS uptake’ and ‘AR selective’ formulated scores ($n = 400$).

Roundworm control practices	SCOPS recommended			AR selective		
	Levels	$n =$	%	Levels	$n =$	%
In the last 12 months how often have you sought advice specifically regarding roundworm control?	At least once	255	64	-	-	-
In the last 12 months how many times have you treated your ewes and lambs for roundworms?	Ewes (< average*)	90	23	Ewes (> average*)	172	43
	Lambs (< average†)	66	17	Lambs (> average†)	195	49
Do you monitor worm egg counts?	Yes	136	34	No	264	66
Do you drench incoming sheep brought onto the farm? ‡	Yes	303	94	No	20	6
Do you withhold sheep from pasture? ‡	Yes	221	68	No	102	32
Have you ever tested for drug resistance?	Yes	51	13	No	349	87
Do you move your animals immediately to clean pasture after treatment?	No	158	40	Yes	244	61
Do you use selective breeding for roundworm control in your flock?	Yes	49	12	-	-	-
Do you graze sheep and cattle together, graze separately or rotate grazing between the two?	Yes – Rotational	84	21	-	-	-
	Yes – Co-graze	134	36	-	-	-

*Ewe treatment average (2) † lamb treatment average (2) ‡ results exclude closed flock farms ($n = 77$)




Fig. 2 - Total number of SCOPS (filled bars) and AR selective (open bars) practices employed by respondents ($n = 400$)

Statistical analysis

Factor analysis

Initial exploratory factor analysis was performed on ordinal (Likert Scale) items related to the general uptake of SCOPS and AR selective practices, in order to identify and evaluate inter-relationships between variables. Based on their covariation, the total number of observed variables was condensed into a smaller set of unobserved (latent) factors. In the development of the proposed models, items within section 2, i.e. general attitudes to roundworm control section were assessed. The procedures for the assessment of factor loadings (correlation coefficients) and reliability analysis (Cronbach alpha) were conducted as described by Hair et al. (2006). Accordingly, based on the study sample size ($n = 400$), in order to achieve

statistical significance for each value with a statistical power of 80 per cent, a minimum threshold of ± 0.30 factor loading was used. Factor loadings below ± 0.30 , or loadings that demonstrated significant loadings across more than one factor i.e. cross loading, were not included within the resultant factor. The internal reliability measure (Cronbach alpha) was set at an approximate minimum threshold of 0.60 with a value > 0.70 indicating a good reliability measure. Factors which demonstrated acceptable factor loadings and Cronbach alpha measures were retained for further analysis. The method of extraction applied was Principal Component Analysis. An orthogonal factor rotation method 'Varimax' was used to interpret the extracted factors.

Structural equation modelling

In order to examine the inter-relationships between the observed and unobserved (latent) variables in the proposed theoretical model (as represented in Figure 1), the analysis was performed using the multivariate analysis technique known as Structural Equation Modelling (SEM). This technique comprises of two parts, the first is the measurement model which represents the relationships between the specified indicators and their latent constructs. The second part is the structural model which then examines the relationships between the model constructs. The relationship between variables as measured by the regression coefficient represents the change in the dependent variable for one unit change in the independent variable. The regression coefficients are standardised (β values) in order to allow direct comparisons of the relative effects of each variable on the dependent variable. The individual effects are estimated independent of the effects of the other variables to allow assessment of individual relationships within the model (i.e. ceteris paribus). All factors were included in both models with the exception of 'Vet service pros' and 'Vet service cons' which were

selected for ‘SCOPS practice uptake’ and ‘AR selective practice uptake’ respectively. The statistical package Lisrel 8.80 (Jöreskog and Sörbom, 2007) was chosen for the purposes of the SEM analysis. Due to the non-normality of the explanatory variable data, a Diagonally Weighted Least Squares (DWLS) method was used to estimate the model parameters. The resulting model output was evaluated for goodness of fit by using the following model fit indices as detailed by Hair et al. (2006); Root Mean Square Error of Approximation (RMSEA), Standardised Root Mean Residual (SRMR), Comparative Fit Index (CFI), Incremental Fit Index (IFI), Goodness of fit (GFI), Adjusted Goodness of Fit Index (AGFI) and Normed Fit Index (NFI).

Results

Participant descriptive statistics

The total number of opt-out letters received from the original 1,930 farmers contacted was 427 (22%), leaving 1,503 farmers eligible to be contacted. The target of 400 completed interviews was achieved with the following numbers of interviews resulting from each region: 65 in the South East, 76 in the South West 74 in Central region, 92 in the North West, 46 in the North East, and 47 in the Islands.

In terms of respondents demographic responses, the majority of farmers (69%) in the survey sample were aged in the 51-65 or >65 year brackets with only 3% of the respondents representing the youngest age bracket (18-35). The number of years earning a living as a farmer was normally distributed with less of a skew towards more experienced farmers. The level of education showed that most respondents (55%) had had some degree of further education, with approximately 35% studying at an agricultural college.

313 In regards to the farming enterprises, almost two thirds of respondents' farms were situated
 314 on either upland or hill grazing land, with over half of the sample population comprising of
 315 mixed livestock farmers and a quarter sheep-only farmers. The proportional flock sizes as
 316 indicated by numbers of breeding ewes are more orientated towards small to medium sized
 317 flocks (i.e. <500 ewes), with a quarter of farms with larger flocks (>500 ewes).

318

319 Table 2

320 Description of latent constructs with corresponding indicators and Cronbach alpha reliability measures (α)

Latent variable	Statement	α	Value and labels	Variable type
Experience	What is your age?	0.764	1 = 18-35; 2 = 36-50; 3 = 51-65; 4 = over 65	Categorical
	How many years have you been earning a living as a farmer?		1 = 10 years or less; 2 = 11-20; 3 = 21-30; 4 = 31-40; 5 = 41-50; 6 = over 51	Categorical
Education	Did you attend a place of further education?	NA	0 = no or yes, education other than agriculture college 1 = yes, Agricultural college	Binary
Ewe numbers	Number of breeding ewes?	NA	1 = 0-100; 2 = 101-200; 3 = 201-500; 4 = 501-1000; 5 = 1001 or more	Categorical
Enterprise type	Is your farm: sheep only, mixed livestock or livestock and arable?	NA	0 = sheep only; 1 = mixed livestock; 2 = livestock and arable	Categorical
Topography	Is your farm designated as lowland, upland or hill?	NA	0 = lowland; 1 = upland; 2 = hill	Ordinal
Worm control knowledge	Knowledge score	NA	0 = none correct; 1 = one correct; 2 = two correct; 3	Ordinal

			= three correct	
Occurrence of worm problems	How would you classify the occurrence of roundworm problems in your flock?	NA	0 = low; 1 = moderate; 2 = high	Ordinal
AR confirmation	Do you have confirmed drug resistance?	NA	1= no 2= yes	Binary
Worm control importance	1. - Roundworm control is important on my farm	0.877	5-point Likert scale ^a	Ordinal
	2. - My roundworm control strategy improves the productiveness of my animals		5-point Likert scale ^a	Ordinal
	3. - Roundworm control is important for the profitability of my farm		5-point Likert scale ^a	Ordinal
	4. - Roundworm control is important for the health & welfare of my animals		5-point Likert scale ^a	Ordinal
AR risk	1. - Wormer resistance is a problem in my region	0.593	5-point Likert scale ^a	Ordinal
	2. - Wormer resistance is a threat to my farming business		5-point Likert scale ^a	Ordinal
Vet service pros	1. - Working with my vet could improve my roundworm control strategy	0.877	5-point Likert scale ^a	Ordinal
	2. - Working out a roundworm control strategy with my vet is cost effective		5-point Likert scale ^a	Ordinal
	3. - Working out a roundworm control strategy with my vet ensures I get reliable advice		5-point Likert scale ^a	Ordinal
Vet service cons	1. - Roundworm control advice provided by vets is too complex	0.81	5-point Likert scale ^a	Ordinal
	2. - Roundworm control advice provided by vets is difficult to implement		5-point Likert scale ^a	Ordinal
SCOPS practice uptake	Number of SCOPS practices implemented	NA	0 = none; 1 = one; 2 = two; 3 = three; 4 = four; 5 =	Ordinal

five; 6 = six; 7 = seven; 8 =
eight; 9 = nine; 10 = ten

AR selective	Number of AR selective practice	NA	0 = none; 1 = one; 2 = two;, Ordinal
practice uptake	implemented		3 = three; 4 = four; 5 =
			five; 6 = six; 7 = seven;

^a 5-point Likert scale: 1 = Strongly disagree; 2 = Disagree; 3 = Unsure; 4 = Agree; 5 = Strongly agree

Results of factor analysis

Both of the models proposed consist of seven single-indicator latent variables and four multiple-indicator latent variables as detailed in Table 2. The exploratory factor analysis established acceptable factor loadings i.e. > 0.70 for all multiple-indicator latent variables (Presented in Appendix A). Additionally, the Cronbach alpha reliability analysis shown in Table 2, demonstrated suitable measures ($\alpha > 0.60$) between all sets of indicators with the exception of 'AR risk'.

Results of structural equation models

Both models reflected a goodness of fit with the observed data as indicated by the following model fit indices as according to Hair et al (2006). Significance was established for all relationships at a 0.05 level, with significant standardised coefficients (total effects) of both models detailed in tables 3 and 4. An illustrated version of the direct influences on SCOPS practice uptake model is presented in Figure 3; however, this was not feasible in the 'AR selective practice' model due to the large number of estimates identified. The SCOPS model fit values were below the maximum threshold of 0.10 for RMSEA at 0.025, and at the 0.08 threshold for SRMR (0.08), for the subsequent fit indices values above 0.90 give an indication of acceptable fit; CFI (0.99), IFI (0.99), GFI (0.98), AGFI (0.97) and NFI (0.96). The SCOPS model explained 54% of the variance in the adoption score of sustainable

parasite control practices. The factors which had the greatest direct positive effects on SCOPS uptake were ‘AR confirmation’ ($\beta = 0.55$) followed by ‘Enterprise type’ ($\beta = 0.30$), ‘AR risk’ ($\beta = 0.21$) and ‘Vet service pro’ ($\beta = 0.20$). The greatest indirect positive influence on SCOPS uptake was ‘Worm control knowledge’ ($\beta = 0.34$) mediated by ‘AR confirmation’ ($\beta = 0.61$). Exogenous factors which were shown to have a positive influence on mediating factors included ‘Ewe numbers’ with a strong effect on ‘AR confirmation’ ($\beta = 0.43$) and a moderate effect on ‘Occurrence of worm problems’ ($\beta = 0.20$). In addition to ‘Education’ with a positive effect on ‘AR risk’ ($\beta = 0.31$) and ‘Worm control importance’ with a positive influence on ‘Vet service pro’ ($\beta = 0.36$). Factors which demonstrated a negative influence on SCOPS uptake through mediating factors included ‘Experience’ on ‘AR risk’ ($\beta = -0.16$) and ‘Worm control knowledge’ ($\beta = -0.31$) as well as ‘Topography’ with moderate influences on ‘Worm control knowledge’ ($\beta = -0.24$).

The AR selective practice model fit indices were as follows; RMSEA (0.050), SRMR (0.083), CFI (0.93), IFI (0.94), GFI (0.97), AGFI (0.96) and NFI (0.90). The AR model explained 47% of the variance in the adoption of recognised AR selective roundworm control practices. Factors shown to have the greatest positive influence on the use of AR selective practices included ‘Vet service con’ with a direct effect on the behavioural outcome ($\beta = 0.14$), in addition to ‘Experience’ ($\beta = 0.12$) and ‘Topography’ ($\beta = 0.08$) which both had indirect influences on AR selective practices. The greatest direct negative influence on AR selective practices was associated with ‘AR confirmation’ ($\beta = -0.67$). Indirect negative influences on AR selective practices included ‘Worm control knowledge’ ($\beta = -0.34$), ‘Ewe numbers’ ($\beta = -0.16$), ‘AR risk’ ($\beta = -0.15$), ‘Education’ ($\beta = -0.11$), ‘Enterprise type’ ($\beta = -0.06$) and ‘Worm control importance’ ($\beta = -0.03$).

The factor ‘AR confirmation’ was shown to be directly influenced positively by ‘Worm control knowledge’ ($\beta = 0.51$), ‘Ewe numbers’ ($\beta = 0.33$) and ‘AR risk’ ($\beta = 0.22$), Indirect

mediated influences included 'Education' ($\beta = 0.16$), 'Enterprise type' ($\beta = 0.09$) and 'Worm control importance' ($\beta = 0.04$). 'AR confirmation' was most negatively influenced by 'Experience' ($\beta = -0.18$) and 'Topography' ($\beta = -0.13$). The factor 'AR risk' attitudes were shown to be most positively influenced directly by 'Occurrence of worm problems' ($\beta = 0.34$), 'Education' ($\beta = 0.26$) and negative influenced by 'Experience' ($\beta = -0.20$) and 'Topography' ($\beta = -0.15$). The factor 'Worm control knowledge' was influenced directly by five factors including most prominently 'Experience', ($\beta = -0.27$) followed by 'Education' ($\beta = 0.21$), 'Topography' ($\beta = -0.18$), 'Enterprise type' ($\beta = 0.17$) and 'Ewe numbers' ($\beta = 0.13$). The factor 'Occurrence of worm problems' was influenced positively by 'Ewe numbers' ($\beta = 0.24$).




Fig.3. SCOPS uptake structural model (standardised solution). Bold arrows represent the direct influences of latent variables on the behavioural latent 'SCOPS practice uptake', with non- bold arrows representing the direct effect influences on other latent variables. The corresponding numbers are the standardised coefficients of the variables in the structural model. Blue variables denote variables that are exogenous i.e. independent from other variables in the model, with green variables taking either exogenous or endogenous roles i.e. influenced by other variables. The orange variable represents the endogenous behavioural latent variable.

400 Table 3.

401 Standardised total effects on SCOPS model latent variables (Standard error values)

Total (direct and indirect) effects on effector variables							
Determinants	‘SCOPS practice uptake’	‘AR Confirmation’	‘Worm control Importance’	‘AR risk’	‘Worm control knowledge’	‘Occurrence of worm problems’	‘Vet service pro’
‘Ewe numbers’	0.25 (0.04)	0.43 (0.08)	-	0.06 (0.03)	-	0.20 (0.03)	-
‘Education’	0.06 (0.03)	-	-	0.31 (0.14)	-	-	-
‘Experience’	-0.14 (0.05)	-0.19 (0.06)	-	-0.16 (0.11)	-0.31 (0.05)	-	-
‘Topography’	-0.08 (0.03)	-0.14 (0.05)	-	-	-0.24 (0.05)	-	-
‘Enterprise type’	0.30 (0.12)	-	-	-	-	-	-
‘AR Confirmation’	0.55 (0.09)	NA	-	-	-	-	-
‘Worm control Importance’	0.07 (0.03)	-	NA	-	-	-	0.36 (0.06)
‘AR risk’	0.21 (0.06)	-	-	NA	-	-	-
‘Worm control knowledge’	0.34 (0.11)	0.61 (0.14)	-	-	NA	-	-
‘Occurrence of worm problems’	0.07 (0.05)	-	-	0.32 (0.18)	-	NA	-
‘Vet service pro’	0.20 (0.07)	-	-	-	-	-	NA
R-square	0.54	0.56	-	0.27	0.16	0.04	0.13

402

403

404

405 Table 4.

406 Standardised total effects on AR model latent variables (Standard error values)

Total (direct and indirect) effects on effector variables							
Determinants	'AR selective practice uptake'	'AR Confirmation'	'Worm control Importance'	'AR risk'	'Worm control knowledge'	'Occurrence of worm problems'	'Vet service e con'
'Ewe numbers'	-0.16 (0.05)	0.33 (0.07)	-	0.08 (0.02)	0.13 (0.03)	0.24 (0.04)	-
'Education'	-0.11 (0.04)	0.16 (0.05)	-	0.26 (0.07)	0.21 (0.06)	-	-
'Experience'	0.12 (0.04)	-0.18 (0.05)	-	-0.20 (0.05)	-0.27 (0.04)	-	-
'Topography'	0.08 (0.04)	-0.13 (0.04)	-	-0.15 (0.05)	-0.18 (0.09)	-	-
'Enterprise type'	-0.06 (0.06)	0.09 (0.07)	-	-	0.17 (0.04)	-	-
'AR Confirmation'	-0.67 (0.10)	NA	-	-	-	-	-
'Worm control Importance'	-0.03 (0.02)	0.04 (0.02)	NA	0.18 (0.05)	-	-	--
'AR risk'	-0.15 (0.08)	0.22 (0.09)	-	NA	-	-	-
'Worm control knowledge'	-0.34 (0.13)	0.51 (0.13)	-	-	NA	-	-
'Occurrence of worm problems'	0.18 (0.15)	0.08 (0.05)	-	0.34 (0.10)	-	NA	-
'Vet service cons'	0.14 (0.07)		-	-	-	-	NA
R-square	0.47	0.44	-	0.28	0.19	0.06	-

407

408 Discussion

409 The results demonstrate that of the nine significant factors positively influencing the uptake
410 of SCOPS recommended practices, the confirmation of AR on a particular holding is shown
411 to have the greatest influence towards the uptake of sustainable parasite control practices.
412 This would suggest that such an event is likely to have the greatest impact on farmer's
413 decision making, which may demonstrate a decisive mechanism for prompting farmers

414 directly affected by AR to assess their treatment efficacies. Farmers may be motivated to
415 modify their parasite control strategies based on the knowledge of which nematode species
416 are resistant to a particular class of anthelmintic, which will help to ensure the preserved
417 effectiveness of the other remaining anthelmintics. The challenge therefore is to encourage
418 farmers to test their treatment efficacies in the absence of indication or a critical event, which
419 has also been acknowledged as a barrier for dairy farmers to reassess their routines regarding
420 mastitis control (Dillon, 2015).

421 The level of farmer's roundworm control knowledge is likely to reflect their awareness and
422 understanding of the topic, which is fundamental to the decision making process. The impact
423 of knowledge on SCOPS uptake emphasises the importance of informing farmers about areas
424 such as roundworm identification, non-chemical control measures and AR as a vital target for
425 influencing farmer's roundworm practices. Furthermore, knowledge was also identified as a
426 strong determinant for establishing AR status which as previously stated may further
427 influence the adoption of SCOPS practices. The negative effect of knowledge on AR practice
428 uptake also demonstrates the influence of SCOPS awareness towards the adoption of
429 sustainable roundworm practices. In another study using SEM, Toma et al (2015) also
430 identified disease control knowledge to directly and indirectly influence farmer's behavioural
431 intentions. The use of farmers' workshops has been one such strategy employed to engage
432 farmers through providing information as well as setting up subsidised faecal egg count
433 monitoring programmes with local veterinary practices during the peak grazing season
434 (Anonymous, 2016). The dual benefits of this type of approach may come from ways of
435 improving motivation as well as providing an added financial incentive. Steers and Porter
436 (1975) suggested motivation may be a result of firstly stimulating an initial interest on a topic
437 (i.e. energising), directing participants to learn and master the topic (director) and then
438 reinforcing the knowledge and skills acquired (i.e. maintenance). The maintenance of

engagement has also been stated as an important aim to achieving behavioural change in the medium to long-term future (Rushmer et al., 2014). The use of economic incentives such as cost-sharing as described in this instance may spur participation from those farmers with a pre-existing interest on the subject, however for those without interest this may have little or no long term effect on the adoption of such sustainable agricultural practices (Rodriguez et al., 2009). The method used to formulate the knowledge score meant that the level of detail in participant's responses was not factored into the analyses. This will therefore have a limiting effect on the depth of understanding attributed to participant responses. Further work may benefit from assessing the influence of superficial vs. in-depth parasite knowledge on the effectiveness of implementing behaviours.

With regards to attitudinal factors, farmers' AR risk perception presented a moderate influence on the uptake of SCOPS practices and a comparable negative influence on AR selective practices. This may suggest that Scottish farmer's perceptions of AR risk in terms of susceptibility and impact may not be as influential as other factors, possibly due to the progressive 'invisible' nature of AR development in comparison with other disease threats (Woodgate and Love, 2012). In fact, the proportion of respondents' disagreeing that AR is a problem in their region or that AR is a threat to their farming business was 42% and 42% respectively. Positive attitudes towards veterinarians' roundworm control services was also shown to influence the uptake of SCOPS practices as would be anticipated due to their prominent role in educating and encouraging sustainable farming practices. The importance of veterinarians as an influential source of roundworm control information was stated by 65% of respondents, and is also widely acknowledged in the literature (Brennan and Christley, 2013; Alarcon et al., 2014). These findings, in support of others e.g. Kaler and Green (2013) reinforce the need to improve interactions between sheep farmers and veterinarians to

463 encourage more farmers to introduce improvements to their current roundworm control
464 strategies, as part of their overall flock health plans.

465 Farming characteristic factors such as ewe numbers and enterprise type were also shown to
466 positively influence SCOPS uptake. The effect of flock size may vary the relative importance
467 and impacts attributed to roundworm control. For instance, larger flocks would typically be
468 more associated with greater stocking densities resulting in a higher parasite infection
469 pressure, due to increased pasture contamination. Hence there is a greater requirement for
470 such farms to employ various measures in order to mitigate production losses, as well as
471 address mounting concerns over reliance on chemical control methods. Willock et al., (1999)
472 also found farm size to be a significant influence to farmer's decision making. Enterprise type
473 was shown to have a considerable direct influence on the uptake of SCOPS practices, which
474 would suggest that farms with a greater diversity of farm enterprises are more likely to adopt
475 'best practice' advice. This would support the findings of other studies where more farm
476 enterprises was shown to influence the adoption of best management practices in cattle
477 production (Kim, 2005). The topography of respondent's farms was also shown to have a
478 relatively small direct influence on adoption of SCOPS behaviours with upland/hill farms less
479 likely to employ such practices. This might be due to the contrasting management systems
480 between lowland and hill farms with greater labour requirements to gather and manage an
481 extensively run flock (Morgan-Davies et al., 2006).

482 Factors that were shown to have a low direct effect on SCOPS uptake included: the
483 occurrence of roundworm control problems, education, topography and perceived
484 roundworm control importance. These factors however demonstrated a greater direct effect
485 through mediating factors such as AR risk, AR confirmation, vet services pro and knowledge.
486 An agricultural college education was shown to positively influence AR risk perception
487 whereas experience was shown to negatively influence numerous factors including

knowledge, AR risk and establishing AR status. The negative influence of other internal factors such as experience suggest that more experienced farmers are less likely to employ sustainable parasite control measures, perhaps due to a greater reliance on their own sense of judgement (Garforth et al., 2013; Kaler and Green, 2013). This concept of self-identity in relation to the importance of farmers own abilities to identify problems poses a likely barrier towards more experienced farmers seeking external guidance regarding roundworm control (Thompson, 2008). This is particularly relevant considering the high proportion of surveyed respondents aged above 51 years of age in contrast with the younger age brackets, which are comparable with most recent agricultural census reports (National Statistics, 2015)

The identification of factors with the greatest influences on best practice uptake can be used to direct future extension programmes towards areas where greatest impact may be expected to occur, such as developing communication strategies highlighting the benefits of diagnostic testing. The utilisation of local veterinary services as a highly trusted resource is likely to appeal most to farmers as this will also facilitate the tailoring of advice to suit the management strategies in their particular enterprises. The main difficulty of this however is the availability of sheep specialist veterinarians with the interest and expertise required to engage farmers on a wider level (Kaler and Green, 2013). Another approach could be to further support the training of animal health advisors as well as those teaching at agricultural colleges, which as demonstrated could help to encourage the next generation of young farmers to adopt best practice parasite management approaches. Finally, lessons could also be taken from other disciplines associated with influencing farmer perceptions and behaviours such as in the agricultural business and marketing sectors. By developing a suite of strategies to address farmers' perceptions and awareness of best practice advice, this will more likely have a greater general impact than using one such approach in isolation.

Conclusions

The use of structural equation modelling has identified a number of significant factors influencing farmer's parasite control behaviours. Both internal and external factors are shown to influence the adoption of SCOPS and AR selective practices including most prominently parasite control knowledge and the identification of AR. Such factors will inform and prompt farmers to think more proactively regarding their roundworm control strategies in order to preserve the effectiveness of remaining anthelmintic treatments. The influence of external factors such as flock size, enterprise type and topography highlight the possible benefits of tailoring future recommendations to suit the range of farming systems present in the sheep farming industry and the challenges associated within these settings.

Acknowledgements

The authors would like to thank all the farmers involved for their active participation in the piloting and running of the telephone surveys. Additional thanks to Fiona Vande Velde for her assistance in the initial stages of the analysis. This study is funded by the RESAS branch of the Scottish Government.

Conflict of interest

None

References

Ajzen, I., 1991. The Theory of Planned Behavior. *Organ Behav Hum Dec* 50, 179-211.

535 Abbott K.A., Taylor M.A., Stubbings L.A., 2012. Sustainable Control of Parasites in Sheep
536 (SCOPS) Sustainable Worm Control Strategies for sheep, 4th Edition, a technical
537 manual for veterinary surgeons and advisors.

538 Alarcon, P., Wieland, B., Mateus, A.L., Dewberry, C., 2014. Pig farmers' perceptions,
539 attitudes, influences and management of information in the decision-making process
540 for disease control. *Prev. Vet. Med.* 116, 223-242.

541 Bartley, D.J., Jackson, E., Johnston, K., Coop, R.L., Mitchell, G.B.B., Sales, J., Jackson, F.,
542 2003. A survey of anthelmintic resistant nematode parasites in Scottish sheep flocks.
543 *Vet. parasitol.* 117, 61-71.

544 Bartley, D.J., Jackson, F., Jackson, E., Sargison, N., 2004. Characterisation of two triple
545 resistant field isolates of *Teladorsagia* from Scottish lowland sheep farms. *Vet.*
546 *parasitol.* 123, 189-199.

547 Bartley, D.J., 2008. Prevalence, characterisation and management of anthelmintic resistance
548 in gastro-intestinal nematodes of Scottish sheep. Unpublished PhD thesis. University
549 of Edinburgh.

550 Brennan, M.L., Christley, R.M., 2013. Cattle producers' perceptions of biosecurity. *BMC*
551 *Vet. Res.* 9, 71.

552 Cawthorne, R.J., Whitehead, J.D., 1983. Isolation of benzimidazole resistant strains of
553 *Ostertagia circumcincta* from British sheep. *Vet. Rec.* 112, 274-277.

554 Coles, G.C., Simkins, K., 1996. Resistance to levamisole. *Vet. Rec.* 139, 124-124.

555 Coles, G.C., 1997. Nematode control practices and anthelmintic resistance on British sheep
556 farms. *Vet. Rec.* 141, 91-93.

557 Coop, R.L., Kyriazakis, I., 2001. Influence of host nutrition on the development and
558 consequences of nematode parasitism in ruminants. *Trends Parasitol* 17, 325-330.

559 Dillon, E.J., 2015. Exploring the role of farmers' attitudes in influencing animal health best
560 practice. Agricultural Economics Society.

561 Edwards-Jones, G., 2006. Modelling farmer decision-making: concepts, progress and
562 challenges. *Anim Sci* 82, 783-790.

563 Garforth, C.J., Bailey, A.P., Tranter, R.B., 2013. Farmers' attitudes to disease risk
564 management in England: a comparative analysis of sheep and pig farmers. *Prev. Vet.
565 Med.* 110, 456-466.

566 Grimshaw, W.T., Hunt, K.R., Hong, C., Coles, G.C., 1994. Detection of anthelmintic
567 resistant nematodes in sheep in southern England by a faecal egg count reduction test.
568 *Vet. Rec.* 135, 372-374.

569 Hair, J.F., Tatham, R.L., Anderson, R.E., Black, W. (2006). *Multivariate Data Analysis*, 6th
570 edition. Prentice Hall.

571 Hong, C., Hunt, K.R., Coles, G.C., 1994. Resistance to Levamisole. *Vet. Rec.* 135, 535-536.

572 Hughes, P.L., Dowling, A.F., Callinan, A.P., 2007. Resistance to macrocyclic lactone
573 anthelmintics and associated risk factors on sheep farms in the lower North Island of
574 New Zealand. *NZ Vet. J.* 55, 177-183.

575 Israel, G. D. (1992). Determining sample size. University of Florida Cooperative Extension
576 Service, Institute of Food and Agriculture Sciences, EDIS.

577 Jackson, F., Coop, R.L., 2000. The development of anthelmintic resistance in sheep
578 nematodes. *J. Parasitol.* 120, S95-S107.

579 Jöreskog, K.G., Sörbom, D., 2007. LISREL8.80: Structural Equation Modeling with the
580 SIMPLIS Command Language. IL Scientific Software International, Chicago, USA.

581 Kaler, J., Green, L.E., 2013. Sheep farmer opinions on the current and future role of
582 veterinarians in flock health management on sheep farms: a qualitative study. *Prev.
583 Vet. Med.* 112, 370-377.

584 Kim, S., Gillespie, J. M., & Paudel, K. P. (2005). The effect of socioeconomic factors on the
585 adoption of best management practices in beef cattle production. *J. Soil Water*
586 *Conserv*, 60, 111-120.

587 Lawrence, K.E., Lethwick, D.M., Rhodes, A.P., Jackson, R., Heuer, C., Pomroy, W.E., West,
588 D.M., Waghorn, T.S., Moffat, J.R., 2007. Management of gastrointestinal nematode
589 parasites on sheep farms in New Zealand. *NZ Vet. J.* 55, 228-234.

590 McMahon, C., McCoy, M., Ellison, S.E., Barley, J.P., Edgar, H.W., Hanna, R.E., Malone,
591 F.E., Brennan, G.P., Fairweather, I., 2013. Anthelmintic resistance in Northern
592 Ireland (III): uptake of 'SCOPS' (Sustainable Control of Parasites in Sheep)
593 recommendations by sheep farmers. *Vet. Parasitol.* 193, 179-184.

594 Mitchell, E.S., Hunt, K.R., Wood, R., McLean, B., 2010. Anthelmintic resistance on sheep
595 farms in Wales. *Vet. Rec.* 166, 650-652.

596 Morgan-Davies, C., Waterhouse, A., Milne, C.E., Stott, A.W., 2006. Farmers' opinions on
597 welfare, health and production practices in extensive hill sheep flocks in Great
598 Britain. *Livest Sci.* 104, 268-277.

599 Morgan, E.R., Hosking, B.C., Burston, S., Carder, K.M., Hyslop, A.C., Pritchard, L.J.,
600 Whitmarsh, A.K., Coles, G.C., 2012. A survey of helminth control practices on sheep
601 farms in Great Britain and Ireland. *Vet. J.* 192, 390-397.

602 National Statistics., 2015. Results from the June 2015 Scottish Agricultural Census. In:
603 *Agriculture, E.a.M. (Ed.) The Scottish Government.*

604 Nieuwhof, G.J., Bishop, S.C., 2005. Costs of the major endemic diseases of sheep in Great
605 Britain and the potential benefits of reduction in disease impact. *Anim Sci* 81, 23-29.

606 Palmer, S., Sully, M., & Fozdar, F., 2009. Farmers, animal disease reporting and the effect of
607 trust: a study of West Australian sheep and cattle farmers. *Rural society* 19, 32-48.

608 Relf, V.E., Morgan, E.R., Hodgkinson, J.E., Matthews, J.B., 2012. A questionnaire study on
609 parasite control practices on UK breeding Thoroughbred studs. *Equine Vet. J.* 44,
610 466-471.

611 Rodriguez, J.M., Molnar, J.J., Fazio, R.A., Sydnor, E., Lowe, M.J., 2009. Barriers to adoption
612 of sustainable agriculture practices: Change agent perspectives. *Renew Agr Food Syst*
613 24, 60-71.

614 Rosenstock, I.M., Strecher, V.J., Becker, M.H., 1988. Social-Learning Theory and the Health
615 Belief Model. *Health Educ Quart* 15, 175-183.

616 Rushmer, R.K., Hunter, D.J., Steven, A., 2014. Using interactive workshops to prompt
617 knowledge exchange: a realist evaluation of a knowledge to action initiative. *Public*
618 *Health* 128, 552-560.

619 Sargison, N.D., Jackson, F., Bartley, D.J., Moir, A.C., 2005. Failure of moxidectin to control
620 benzimidazole-, levamisole- and ivermectin-resistant *Trichostrongylus axei* in a
621 sheep flock. *Vet. Rec.* 156, 105-109.

622 Sargison, N.D., Jackson, F., Bartley, D.J., Wilson, D.J., Stenhouse, L.J., Penny, C.D., 2007.
623 Observations on the emergence of multiple anthelmintic resistance in sheep flocks in
624 the south-east of Scotland. *Vet. Parasitol.* 145, 65-76.

625 Steers, R.M., Porter, L.W., 1975. *Motivation and work behavior*. New York: McGraw-Hill.

626 Suter, R.J., McKinnon, E.J., Perkins, N.R., Besier, R.B., 2005. The effective life of
627 ivermectin on Western Australian sheep farms--a survival analysis. *Prev. Vet. Med.*
628 72, 311-322.

629 Sutherland, I.A., Lee, D.L., Lewis, D., 1988. Detection of benzimidazole resistance in
630 trichostrongylid nematodes. *Parasitol. today* 4, 22-24.

631 Thomas, E., Morgan, E., Paton N., 2015. WAARD project: Wales Against Anthelmintic
 632 Resistance. In: Wales, M.P. (Ed.). Last accessed 12MAY16.
 633 http://hccmpw.org.uk/farming/animal_health_and_welfare/anthelmintic_resistance/
 634 Thompson, L.J., 2008. Extending Complex Agricultural Research in Australia: The case of
 635 integrated parasite management in sheep. Institute for Rural Futures. University of
 636 New England, Armidale, Australia.
 637 Toma, L., Low, J.C., Vosough Ahmadi, B., Matthews, L., Stott, A.W., 2015. An analysis of
 638 cattle farmers' perceptions of drivers and barriers to on-farm control of *Escherichia*
 639 *coli* O157. *Epidemiol. Infect.* 143, 2355-2366.
 640 Toma, L., Stott, A.W., Heffernan, C., Ringrose, S., Gunn, G.J., 2013. Determinants of
 641 biosecurity behaviour of British cattle and sheep farmers-a behavioural economics
 642 analysis. *Prev. Vet. Med.* 108, 321-333.
 643 Vande Velde, F., Claerebout, E., Cauberghe, V., Hudders, L., Van Loo, H., Vercruysse, J.,
 644 Charlier, J., 2015. Diagnosis before treatment: Identifying dairy farmers' determinants
 645 for the adoption of sustainable practices in gastrointestinal nematode control. *Vet.*
 646 *Parasitol.* 212, 308-317.
 647 Wauters, E., Rojo-Gimeno, C., 2014. Socio-psychological veterinary epidemiology. A new
 648 discipline for an old problem. Society for Veterinary Epidemiology and Preventive
 649 Medicine (SVEPM) Conference, Dublin, Ireland, 26th to 28th March
 650 Willock, J., Deary, I.J., Edwards-Jones, G., Gibson, G.J., McGregor, M.J., Sutherland, A.,
 651 Dent, J.B., Morgan, O., Grieve, R., 1999. The role of attitudes and objectives in
 652 farmer decision making: Business and environmentally-oriented behaviour in
 653 Scotland. *J Agr Econ* 50, 286-303.
 654 Woodgate, R.G., Love, S., 2012. WormKill to WormBoss--can we sell sustainable sheep
 655 worm control? *Vet. Parasitol.* 186, 51-57.

656 Yamane, T., 1967. Statistics, An Introductory Analysis, 2nd Ed., New York: Harper and
657 Row.

658 **Website references**

659 Anonymous., 2013. <http://www.nationalsheep.org.uk/news/1241/>. Last accessed 12MAY16.

660 Anonymous., 2016a. <http://www.wormboss.com.au/>. Last accessed 12MAY16.

661 Anonymous., 2016b. [http://gwlad.gov.wales/famingconnect/130614-faecal-egg-](http://gwlad.gov.wales/famingconnect/130614-faecal-egg-counting/?jsessionid=D091BF98ED0C3F35DDEC77111011CCA3?lang=en&skip=)
662 [counting/?jsessionid=D091BF98ED0C3F35DDEC77111011CCA3?lang=en&skip=](http://gwlad.gov.wales/famingconnect/130614-faecal-egg-counting/?jsessionid=D091BF98ED0C3F35DDEC77111011CCA3?lang=en&skip=).
663 Last Accessed 12MAY16.

664

665